

AGE		VACCINE	✓
BIRTH	Opt 1	Oral Polio Tuberculosis Hepatitis B*	
	Opt 2		
VISIT 1	6 weeks	Oral Polio Diphtheria Tetanus Pertussis (acellular) Haemophilus influenzae type B (Hib) Hepatitis B (Hb) Polio (Inactive Polio Vaccine)	
	8 weeks		
VISIT 2	10 weeks	Rotavirus** Pneumococcal Diphtheria Tetanus Pertussis (acellular) Haemophilus influenzae type B (Hib) Hepatitis B (Hb) Polio (Inactivated Polio Vaccine)	
	12 weeks		
VISIT 3	14 weeks	Pneumococcal*** Diphtheria Tetanus Pertussis (acellular) Haemophilus influenzae type B (Hib) Hepatitis B (Hb) Polio (Inactivated Polio Vaccine)	
	16 weeks		
9 months		Measles Chickenpox Pneumococcal***	
From 12 months		Hepatitis A Pneumococcal (If private schedule option is followed)	
15 months		Measles, Mumps, Rubella (MMR)	
18 months		Diphtheria Tetanus Pertussis (acellular) Haemophilus influenzae type B (Hib) Hepatitis B (Hb) Polio (Inactivated Polio Vaccine)	
		Hepatitis A	
4 - 6 years		Diphtheria Tetanus Pertussis (acellular)	
		Measles, Mumps, Rubella (MMR)	
Females 10 years +		Cervical Cancer (3 Doses; 0,1,6 months)	
12 years		Tetanus, diphtheria	

information

vaccination

www.Snappi.co.za



\* If a hepatitis B vaccine at birth, a six-in-one vaccination can be given 6, 10, 14 weeks

\*\* Rotavirus vaccine should not be given after 24 weeks

\*\*\* If DOH Schedule is followed Pneumococcal vaccine not given here. 3rd dose given at 9 months

**SNAPPY**®